

POSTER PRESENTATION

Open Access

Serial T2-mapping to quantitatively monitor resorption of myocardial edema following acute myocardial infarction

Gunnar Lund^{1*}, Kai Muellerleile², Peter Bannas¹, Julia Cuerlis¹, Dominik Barz¹, Ulf K Radunski², Christian Stehning³, Bernhard Schnackenburg³, Karsten Sydow², Gerhard Adam¹

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Currently, myocardial edema monitoring after acute myocardial infarction (AMI) is based on visualization of the region with increased signal-intensity on T2-weighted images. T2-mapping is a promising novel cardiac magnetic resonance imaging (CMR) technique to quantitatively assess myocardial edema. The purpose of the study was to quantitatively evaluate resorption of myocardial edema following AMI using T2-mapping.

Methods

CMR (1.5 Tesla Philips Achieva) was performed in 20 patients within seven days after a reperfused AMI (Baseline) and at one, three and six months follow-up, respectively. A free-breathing, navigator-gated multi-echo sequence was used for T2-mapping. T2-maps were calculated from fifteen echoes using a dedicated plug-in written for OsiriX software. Serial T2 values were assessed using a six-segment

model (Figure). Infarcted and remote segments were defined by using information from corresponding late-enhancement images (Figure).

Results

A significant decrease of the T2 time was found in infarcted segments from baseline to one month follow-up (82 ± 19 vs. 70 ± 7 ms; $p < 0.05$), but not between one and three (70 ± 7 vs. 62 ± 8 ; $p = \text{ns}$) or three and six (62 ± 8 vs. 62 ± 5 ms; $p = \text{ns}$) month follow-up. Identical T2 times were found in remote segments at baseline, one, three and six month follow-up (61 ± 6 vs. 61 ± 5 vs. 59 ± 7 vs. 58 ± 2 ms; $p = \text{ns}$). The T2_{infarct}/T2_{remote} ratio decreased from baseline to one month follow-up (1.34 ± 0.26 vs. 1.15 ± 0.10 ; $p < 0.05$) and from one to three month follow-up (1.15 ± 0.10 vs. 1.04 ± 0.06 ; $p = 0.05$). No significant change was found for the T2_{infarct}/T2_{remote} ratio from three to six month follow-up (1.04 ± 0.06 vs. 1.07 ± 0.09 ; $p = \text{ns}$).




Figure 1 CMR of a patient with acute posterior infarction. The infarct and the myocardial edema is seen on the DE- and the T2w-CMR, respectively. The T2-Map shows increased T2 time in the infarcted area. A segmental evaluation was performed in six segments.

¹Radiology, University Hospital, Hamburg, Germany
Full list of author information is available at the end of the article

Conclusions

Serial T2-mapping enables monitoring of edema resorption following acute myocardial infarction. The magnitude of edema resorption occurs in the first month after AMI and the T2 values normalize within 3 months after AMI. Thereafter, no further T2 time reduction is observed.

Funding

None

Author details

¹Radiology, University Hospital, Hamburg, Germany. ²Cardiology, University Hospital, Hamburg, Germany. ³Philips Research, Hamburg, Germany.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-P179

Cite this article as: Lund *et al.*: Serial T2-mapping to quantitatively monitor resorption of myocardial edema following acute myocardial infarction. *Journal of Cardiovascular Magnetic Resonance* 2013 **15**(Suppl 1):P179.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

